

USE OF IMMUNOHISTOCHEMISTRY AS AN ADJUNCT IN THE DIAGNOSIS OF LIMITED ADENOCARCINOMA OF THE PROSTATE CANCER

Jonathan Epstein

- The use of immunohistochemistry for basal cell markers and AMACR for the diagnosis of limited adenocarcinoma of the prostate should be used as an adjunct to the H&E diagnosis as there is false positive and negative staining with these markers.
- Examples of labeling of prostate adenocarcinoma with basal cell markers include: aberrant scattered HMWCK staining of cancer cells; rarely the retention of a basal cell layer in carcinoma; and diffuse expression of p63 in some prostate cancers.
- Basal cell markers are useful in distinguishing mimickers, such as adenosis, atrophy, radiation changes, and sclerosing adenosis, from prostatic adenocarcinoma recognizing that in some cases the staining pattern between mimickers and prostate carcinoma overlap.
- Basal cell stains are helpful but are associated with pitfalls in the diagnosis of HGPIN and its distinction from carcinoma, PIN-like ductal adenocarcinoma, and intraductal carcinoma.
- Although the vast majority of prostate adenocarcinomas label with antisera to PSA, there are some that are negative where the P501S (prostein) and PSMA (prostate specific membrane antigen) may be positive.
- Other examples of useful immunohistochemistry include stains for CD68 to distinguish prostatic xanthoma and nonspecific granulomatous prostatitis from high grade prostate cancer.

Use of Basal Cell Markers and AMACR to Diagnose Limited Carcinoma

The most commonly used antibody to label basal cells in benign mimickers of prostate cancer, is high molecular weight cytokeratin (34BE12, cytokeratin 5/6). High molecular weight cytokeratin immunoreactivity in benign glands is localized to the cytoplasm of basal cells and is negative in prostate cancer. More recently, antibodies to p63 have been shown to label the nuclei of basal cells in benign prostatic lesions.

Several studies comparing high molecular weight cytokeratin and p63 have showed p63 to be slightly superior. One study demonstrated that ck5/6 was superior to 34BE12, although only a minority of pathologists use ck5/6. The use of a double cocktail combining HMWCK and p63 can increase the sensitivity of basal cell detection with a decrease in staining variability.

The use of high molecular weight cytokeratin or p63 in a focus with only a few atypical glands is not as diagnostic, since benign glands may not show uniform positivity with these markers. Negative staining for basal cell markers is most diagnostic when more than a few glands are present for evaluation and the morphologic features are very suspicious for carcinoma. Rather than used to establish a diagnosis of cancer, we use these antibodies to help verify a suspicious focus as cancer. If we favor, although are not

sure, that a focus is benign and the basal cell stains are negative, we will diagnose it as atypical rather than as cancer. In a small focus of atypical glands on prostate biopsy, negative staining for high molecular weight cytokeratin should not necessarily lead to a definitive malignant diagnosis in all cases, as almost half these biopsies on follow-up sampling are benign. If we are confident the focus is benign and stains performed at an outside institution are negative in a small focus of glands, we will still diagnose the focus as benign since certain mimickers of prostate cancer may not react with these antibodies

Alpha-methylacyl-CoA-racemase (AMACR), an enzyme involved in the beta-oxidation of branched-chain fatty acids, is significantly up-regulated in prostate cancer. Antibodies have been developed against its gene product, P504S protein. By immunohistochemistry, the majority of prostate cancers are positive for AMACR, the sensitivity varying amongst studies from 82%-100%. Often the staining is fine dot-like and luminal. Although the data is somewhat conflicting, some studies have shown relative decrease AMACR immunoreactivity in foamy gland, atrophic, and pseudohyperplastic prostate cancers. AMACR staining of PIN and mimickers of prostate cancer is discussed in chapters 5 and 7, respectively. As negative staining for basal cell markers especially in a small focus of atypical glands is not necessarily diagnostic of prostate cancer, positive staining for AMACR can increase the level of confidence in establishing a definitive malignant diagnosis.

Different cocktails have been investigated combining antibodies for AMACR and basal cell specific markers. One combination is with antibodies to p63 which label basal cell nuclei of benign glands and AMACR which stains cytoplasm of cancer. Although these authors have reported that this cocktail is essentially equal to each antibody used separately, in our experience a problem with this cocktail is that in some cases stains for p63 show some background staining of the cytoplasm in benign glands, which can be confused with AMACR immunoreactivity. With small foci of atypical glands, the lesion may not survive sectioning to do separate stains for basal cell markers and AMACR on different slides. A triple stain cocktail using a brown chromogen for both high molecular weight cytokeratin and p63 and a red chromogen for AMACR optimizes the preservation of tissue for immunohistochemistry and has been shown to be better than basal cell markers by themselves.

Hedrick L, Epstein JI. Use of keratin 903 as an adjunct in the diagnosis of prostate carcinoma. *Am J Surg Pathol* 1989;13:389-96.

O'Malley FP, Grignon DJ, Shum DT. Usefulness of immunoperoxidase staining with high-molecular-weight cytokeratin in the differential diagnosis of small-acinar lesions of the prostate gland. *Virchows Arch A Pathol Anat Histopathol* 1990;417:191-6.

Shah IA, Schlageter MO, Stinnett P, Lechago J. Cytokeratin immunohistochemistry as a diagnostic tool for distinguishing malignant from benign epithelial lesions of the prostate. *Mod Pathol* 1991;4:220-4.

Wojno KJ, Epstein JI. The utility of basal cell-specific anti-cytokeratin antibody (34 beta E12) in the diagnosis of prostate cancer. A review of 228 cases. *Am J Surg Pathol* 1995;19:251-60.

Shah RB, Zhou M, LeBlanc M, Snyder M, Rubin MA. Comparison of the basal cell-specific markers, 34betaE12 and p63, in the diagnosis of prostate cancer. *Am J Surg Pathol* 2002;26:1161-8.

Weinstein MH, Signoretti S, Loda M. Diagnostic utility of immunohistochemical staining for p63, a sensitive marker of prostatic basal cells. *Mod Pathol* 2002;15:1302-8.

Parsons JK, Gage WR, Nelson WG, De Marzo AM. p63 protein expression is rare in prostate adenocarcinoma: implications for cancer diagnosis and carcinogenesis. *Urology* 2001;58:619-24.

Abrahams NA, Ormsby AH, Brainard J. Validation of cytokeratin 5/6 as an effective substitute for keratin 903 in the differentiation of benign from malignant glands in prostate needle biopsies. *Histopathology* 2002;41:35-41.

Rubin MA, Zhou M, Dhanasekaran SM, et al. alpha-Methylacyl coenzyme A racemase as a tissue biomarker for prostate cancer. *JAMA* 2002;287:1662-70.

Shah RB, Kunju LP, Shen R, LeBlanc M, Zhou M, Rubin MA. Usefulness of basal cell cocktail (34betaE12 + p63) in the diagnosis of atypical prostate glandular proliferations. *Am J Clin Pathol* 2004;122:517-23.

Zhou M, Shah R, Shen R, Rubin MA. Basal cell cocktail (34betaE12 + p63) improves the detection of prostate basal cells. *Am J Surg Pathol* 2003;27:365-71.

Halushka MK, Kahane H, Epstein JI. Negative 34betaE12 staining in a small focus of atypical glands on prostate needle biopsy: a follow-up study of 332 cases. *Hum Pathol* 2004;35:43-6.

Zhou M, Chinnaiyan AM, Kleer CG, Lucas PC, Rubin MA. Alpha-Methylacyl-CoA racemase: a novel tumor marker over-expressed in several human cancers and their precursor lesions. *Am J Surg Pathol* 2002;26:926-31.

Luo J, Zha S, Gage WR, et al. Alpha-methylacyl-CoA racemase: a new molecular marker for prostate cancer. *Cancer Res* 2002;62:2220-6.

Jiang Z, Woda BA, Rock KL, et al. P504S: a new molecular marker for the detection of prostate carcinoma. *Am J Surg Pathol* 2001;25:1397-404.

Jiang Z, Wu CL, Woda BA, et al. P504S/alpha-methylacyl-CoA racemase: a useful marker for diagnosis of small foci of prostatic carcinoma on needle biopsy. *Am J Surg Pathol* 2002;26:1169-74.

Jiang Z, Wu CL, Woda BA, et al. Alpha-methylacyl-CoA racemase: a multi-institutional study of a new prostate cancer marker. *Histopathology* 2004;45:218-25.

Beach R, Gown AM, De Peralta-Venturina MN, et al. P504S immunohistochemical detection in 405 prostatic specimens including 376 18-gauge needle biopsies. *Am J Surg Pathol* 2002;26:1588-96.

Magi-Galluzzi C, Luo J, Isaacs WB, Hicks JL, de Marzo AM, Epstein JI. Alpha-methylacyl-CoA racemase: a variably sensitive immunohistochemical marker for the diagnosis of small prostate cancer foci on needle biopsy. *Am J Surg Pathol* 2003;27:1128-33.

Zhou M, Jiang Z, Epstein JI. Expression and diagnostic utility of alpha-methylacyl-CoA-racemase (P504S) in foamy gland and pseudohyperplastic prostate cancer. *Am J Surg Pathol* 2003;27:772-8.

Kunju LP, Rubin MA, Chinnaiyan AM, Shah RB. Diagnostic usefulness of monoclonal antibody P504S in the workup of atypical prostatic glandular proliferations. *Am J Clin Pathol* 2003;120:737-45.

Farinola MA, Epstein JI. Utility of immunohistochemistry for alpha-methylacyl-CoA racemase in distinguishing atrophic prostate cancer from benign atrophy. *Hum Pathol* 2004;35:1272-8.

Jiang Z, Iczkowski KA, Woda BA, Tretiakova M, Yang XJ. P504S immunostaining boosts diagnostic resolution of "suspicious" foci in prostatic needle biopsy specimens. *Am J Clin Pathol* 2004;121:99-107.

Zhou M, Aydin H, Kanane H, Epstein JI. How often does alpha-methylacyl-CoA-racemase contribute to resolving an atypical diagnosis on prostate needle biopsy beyond that provided by basal cell markers? *Am J Surg Pathol* 2004;28:239-43.

Kunju LP, Chinnaiyan AM, Shah RB. Comparison of monoclonal antibody (P504S) and polyclonal antibody to alpha methylacyl-CoA racemase (AMACR) in the work-up of prostate cancer. *Histopathology* 2005;47:587-96.

Molinie V, Fromont G, Sibony M, et al. Diagnostic utility of a p63/alpha-methyl-CoA-racemase (p504s) cocktail in atypical foci in the prostate. *Mod Pathol* 2004;17:1180-90.

Hameed O, Sublett J, Humphrey PA. Immunohistochemical stains for p63 and alpha-methylacyl-CoA racemase, versus a cocktail comprising both, in the diagnosis of prostatic carcinoma: a comparison of the immunohistochemical staining of 430 foci in radical prostatectomy and needle biopsy tissues. *Am J Surg Pathol* 2005;29:579-87.

Sanderson SO, Sebo TJ, Murphy LM, Neumann R, Slezak J, Cheville JC. An analysis of the p63/alpha-methylacyl coenzyme A racemase immunohistochemical cocktail stain in prostate needle biopsy specimens and tissue microarrays. *Am J Clin Pathol* 2004;121:220-5.

Browne TJ, Hirsch MS, Brodsky G, Welch WR, Loda MF, Rubin MA. Prospective evaluation of AMACR (P504S) and basal cell markers in the assessment of routine prostate needle biopsy specimens. *Hum Pathol* 2004;35:1462-8.

Jiang Z, Li C, Fischer A, Dresser K, Woda BA. Using an AMACR (P504S)/34betaE12/p63 cocktail for the detection of small focal prostate carcinoma in needle biopsy specimens. *Am J Clin Pathol* 2005;123:231-6.

Hameed O, Humphrey PA. p63/AMACR antibody cocktail retaining of prostate needle biopsy tissues after transfer to charged slides: a viable approach in the diagnosis of small atypical foci that are lost on block sectioning. *Am J Clin Pathol* 2005;124:708-15.

Use of Basal Cell Markers in Radiated Prostate

In addition to radical prostatectomy, external beam radiation and or interstitial radiotherapy (brachytherapy) are currently among the most common options available for the management of localized prostate cancer with a curative intent. Within the nonneoplastic prostatic glands, radiation results in glandular atrophy, squamous metaplasia, and cytologic atypia. Though one may find vascular radiation changes and stromal fibrosis, the stromal atypia characteristic of radiation in other organs is not usually seen. The degree of cytologic atypia in non neoplastic glands and degree of stromal fibrosis appear to be higher after brachytherapy compared to external beam radiation. Furthermore, the marked epithelial atypia tend to persist for a longer time (up to 6 years) following brachytherapy.

The distinction between irradiated nonneoplastic prostatic glands and carcinoma is best made on the architectural pattern of the glands. Within the radiated normal prostate, glands maintain their normal architectural configuration. In contrast to carcinoma, the nonneoplastic glands are separated by a modest amount of prostatic stroma. On higher magnification, there is piling up of the nuclei within irradiated normal prostate as well as an occasional recognizable basal cell layer. Multilayered cells in radiated benign glands frequently appear slightly spindled resembling urothelial metaplasia. The finding of scattered markedly atypical nuclei within well-formed acini is typical of radiated benign glands and rare in prostate carcinoma. Prostate carcinomas that are sufficiently differentiated to form glands rarely manifest the degree of atypia seen with radiation, and if present would be more uniformly present in all cells. Radiated nuclei showing atypia also have a degenerative, hyperchromatic smudgy appearance as opposed to malignant prostatic nuclei that usually contain prominent nucleoli, although occasional nucleoli can be seen in benign prostate glands with radiation affect. Irradiated nonneoplastic glands often are atrophic, in contrast to gland-forming prostatic adenocarcinomas that typically have abundant cytoplasm.

Radiated adenocarcinoma of the prostate may show either no recognizable difference from nonradiated cancer or the effects of radiation damage. In order to diagnose either pattern of cancer, the key feature is that architecturally the findings are inconsistent with benign glands. The presence of closely packed glands with a haphazard

infiltrative growth pattern is typical of adenocarcinoma without treatment affect and cannot be attributed to radiation change. Similarly, the presence of numerous infiltrating individual epithelial cells is diagnostic of carcinoma with treatment affect. Cancers not showing any treatment effect have typical prostate cancer nuclei with prominent nucleoli and glands with a modest amount of cytoplasm. Cancers with radiation effect typically demonstrate individual cells with abundant vacuolated cytoplasm or single cells with indistinct cytoplasm. Nuclei lack apparent nucleoli and are either large with bizarre shapes or pyknotic with smudged chromatin.

It has been demonstrated that high molecular weight cytokeratin immunohistochemistry can aid in the diagnosis of irradiated prostate by identifying basal cells within benign radiated glands. Expression of alpha-methylacyl-coenzyme A racemase (P504S) is usually maintained in radiated adenocarcinoma.

Bostwick DG, Egbert BM, Fajardo LF. Radiation injury of the normal and neoplastic prostate. *Am J Surg Pathol* 1982;6:541-51.

Magi-Galluzzi C, Sanderson H, Epstein JI. Atypia in nonneoplastic prostate glands after radiotherapy for prostate cancer: duration of atypia and relation to type of radiotherapy. *Am J Surg Pathol* 2003;27:206-12.

Brawer MK, Nagle RB, Pitts W, Freiha F, Gamble SL. Keratin immunoreactivity as an aid to the diagnosis of persistent adenocarcinoma in irradiated human prostates. *Cancer* 1989;63:454-60.

Yang XJ, Laven B, Tretiakova M, et al. Detection of alpha-methylacyl-coenzyme A racemase in postradiation prostatic adenocarcinoma. *Urology* 2003;62:282-6.

Crook JM, Bahadur YA, Robertson SJ, Perry GA, Esche BA. Evaluation of radiation effect, tumor differentiation, and prostate specific antigen staining in sequential prostate biopsies after external beam radiotherapy for patients with prostate carcinoma. *Cancer* 1997;79:81-9.

Martens MB, Keller JH. Routine immunohistochemical staining for high-molecular weight cytokeratin 34-beta and alpha-methylacyl CoA racemase (P504S) in postirradiation prostate biopsies. *Mod Pathol* 2006;19:287-90.

Immunohistochemistry for Selected Mimickers of Prostate Adenocarcinoma

Adenosis

Adenosis

Lobular

Cancer

Haphazard growth pattern

Small glands share features with admixed larger glands	Small glands differ from adjacent benign glands
Pale-clear cytoplasm	Occasionally amphophilic cytoplasm
Medium sized nucleoli	Occasionally large nucleoli
Blue mucinous secretions rare	Blue mucinous secretions common
Corpora amylacea common	Corpora amylacea rare
Basal cells present	Basal cells absent

Features Shared in Adenosis and Cancer

- Crowded glands
- Crystalloids
- Medium sized nucleoli
- Scattered poorly formed glands and singles
- Minimal infiltration at periphery
- AMACR immunoreactivity

Atrophy

Post-atrophic hyperplasia (PAH) also often appears basophilic at low power. It consists of acini that are small and mostly round that are arranged in a lobular distribution. Often these acini appear to be surrounding a somewhat dilated “feeder” duct. Many of these lesions frequently resemble normal appearing resting breast lobules, and are referred to by some authors as lobular atrophy. The lesions appear hyperplastic since the close packing of multiple small acini suggests that there is an increase in their number compared to normal tissue. PAH glands have a much higher proliferation rate than nonatrophic benign glands. Although the glands may appear infiltrative, they appear invasive as a patch not as individual glands infiltrating in between larger benign glands. The basophilic appearance of glands of atrophy is due to their scant cytoplasm and crowded nuclei such that at low magnification one is merely seeing a nuclear outline of the gland.

When there are concerns as to whether a focus represents PAH or adenocarcinoma, immunohistochemistry with antibodies to high molecular weight cytokeratin or p63 can be performed to resolve the issue, as PAH uniformly labels with basal cell markers. As opposed to partial atrophy (see below), PAH uncommonly expresses racemase.

Partial atrophy, another variant of atrophy, is the most common mimicker of prostate. Partial atrophy may still retain the lobular pattern of PAH, or have more of a disorganized diffuse appearance. Partial atrophy lacks the basophilic appearance of fully developed atrophy (simple atrophy, PAH) as the nuclei are more spaced apart. The presence of crowded glands with pale cytoplasm may lead to an overdiagnosis of low-grade adenocarcinoma. At higher power, however, the glands have benign features characterized by undulating luminal surfaces with papillary infolding. Most carcinomas have more straight, even luminal borders. In addition, the glands are partially atrophic with nuclei in areas reaching the full height of the cytoplasm. The nuclear features in partial atrophy tend to be relatively benign without prominent nucleoli, although nuclei may appear slightly enlarged with small nucleoli. As with adenosis, partial atrophy typically has a patchy basal cell layer and may express racemase.

Sclerosing Adenosis

Adenocarcinomas of the prostate composed of an admixture of glands, poorly formed glandular structures, and single cells would be assigned a high Gleason score (7 or 8). Prostatic adenocarcinomas with these scores are only rarely seen as limited foci within a TURP. The finding of only one or several small foci of a cellular lesion suspicious for high-grade carcinoma should prompt a consideration of sclerosing adenosis. Furthermore, although sclerosing adenosis may be minimally infiltrative at its perimeter, the lesion is still relatively circumscribed in contrast to high-grade prostate adenocarcinoma.

The glandular structures in sclerosing adenosis resemble those seen in ordinary adenosis. They are composed of cells with pale to clear cytoplasm and relatively benign-appearing nuclei. In many of the glandular structures, a basal cell layer can be identified on H&E-stained sections. This contrasts to carcinoma, where basal cells are absent. Sclerosing adenosis contains a dense spindle cell component that is typically lacking in adenocarcinomas. Usually, adenocarcinomas of the prostate show no apparent stromal response or at most a hypocellular fibrotic reaction. A rather unique feature of sclerosing adenosis is the presence of a hyaline sheath-like structure around some of the glands. The glands in ordinary adenocarcinoma lack such a collarette and have a “naked” appearance as they infiltrate the stroma.

Sclerosing adenosis contains a basal cell layer around most of the glandular structures as well as among the individual cells and cords of cells. The basal cells within sclerosing adenosis, however, are distinctive in their immunophenotypical staining and differ from ordinary basal cells. Ordinary basal cells of the prostate show no myoepithelial cell differentiation. They lack staining for muscle specific actin and ultrastructurally do not show contractile elements. Within sclerosing adenosis, the basal cells show muscle specific actin positivity consistent with myoepithelial cell differentiation. The dense spindle cell component in sclerosing adenosis also shows partial staining with keratin and muscle-specific actin consistent with myoepithelial cell

differentiation. Ultrastructural examination of several of these cases has verified their myoepithelial differentiation. There is no known association between sclerosing adenosis and adenocarcinoma of the prostate.

Hedrick L, Epstein JI. Use of keratin 903 as an adjunct in the diagnosis of prostate carcinoma. *Am J Surg Pathol* 1989;13:389-96.

Yang XJ, Wu CL, Woda BA, et al. Expression of alpha-Methylacyl-CoA racemase (P504S) in atypical adenomatous hyperplasia of the prostate. *Am J Surg Pathol* 2002;26:921-5.

Amin MB, Tamboli P, Varma M, Srigley JR. Postatrophic hyperplasia of the prostate gland: a detailed analysis of its morphology in needle biopsy specimens. *Am J Surg Pathol* 1999;23:925-31.

Ruska KM, Sauvageot J, Epstein JI. Histology and cellular kinetics of prostatic atrophy. *Am J Surg Pathol* 1998;22:1073-7.

Beach R, Gown AM, De Peralta-Venturina MN, et al. P504S immunohistochemical detection in 405 prostatic specimens including 376 18-gauge needle biopsies. *Am J Surg Pathol* 2002;26:1588-96.

Farinola MA, Epstein JI. Utility of immunohistochemistry for alpha-methylacyl-CoA racemase in distinguishing atrophic prostate cancer from benign atrophy. *Hum Pathol* 2004;35:1272-8.

Oppenheimer JR, Wills ML, Epstein JI. Partial atrophy in prostate needle cores: another diagnostic pitfall for the surgical pathologist. *Am J Surg Pathol* 1998;22:440-5.

Grignon DJ, Ro JY, Srigley JR, Troncoso P, Raymond AK, Ayala AG. Sclerosing adenosis of the prostate gland. A lesion showing myoepithelial differentiation. *Am J Surg Pathol* 1992;16:383-91.

Jones EC, Clement PB, Young RH. Sclerosing adenosis of the prostate gland. A clinicopathological and immunohistochemical study of 11 cases. *Am J Surg Pathol* 1991;15:1171-80.

Sakamoto N, Tsuneyoshi M, Enjoji M. Sclerosing adenosis of the prostate. Histopathologic and immunohistochemical analysis. *Am J Surg Pathol* 1991;15:660-7.

Luque RJ, Lopez-Beltran A, Perez-Seoane C, Suzigan S. Sclerosing adenosis of the prostate. Histologic features in needle biopsy specimens. *Arch Pathol Lab Med* 2003;127:e14-6.

Pitfalls with Basal Cell Markers:
Aberrant Staining; Cancers with Retention of Basal Cells; p63+ Cancer

Uncommonly, one can see occasional cancer cells that are positive for antibodies to high molecular weight cytokeratin and less likely p63, yet as long as these cells are not in a basal cell distribution, these cells represent aberrant expression of the antigen in cancer.

Rare lesions with the appearance of prostate cancer show high molecular weight cytokeratin staining in a basal cell distribution either from retention of basal cells by early invasive cancer or from high grade PIN outpouching. The lack of adjacent PIN in some cases and the large ratio of small atypical glands to PIN glands argue against high grade PIN outpouching as the sole explanation. In cases with adjacent high grade PIN, a comparison of the proximity and number of the small, atypical, infiltrative appearing glands to high grade PIN is helpful. The diagnosis of prostate cancer in the face of positive high molecular weight cytokeratin basal cell staining should be made with extreme caution, only in the face of unequivocal cancer on the H&E stain.

There are also uncommon prostate adenocarcinomas that diffusely express p63, yet not HMWCK. Many of these tumors have distinctive morphology composed of atrophic glands lined by hyperchromatic nuclei that are often slightly spindled and minimally multilayered.

Oliai BR, Kahane H, Epstein JI. Can basal cells be seen in adenocarcinoma of the prostate?: an immunohistochemical study using high molecular weight cytokeratin (clone 34betaE12) antibody. *Am J Surg Pathol* 2002;26:1151-60.

Ali T, Epstein JI. False positive labeling of prostate cancer with high molecular weight cytokeratin: p63 a more specific immunomarker for basal cells. *Am J Surg Pathol* (December) 32:1890-5, 2008.

Osunkoya AO, Hansel DE, Sun X, Netto GJ, Epstein JI. Aberrant diffuse expression of p63 in adenocarcinoma of the prostate on needle biopsy and radical prostatectomy: report of 21 cases. *Am J Surg Pathol*. 2008 Mar;32(3):461-7.

Use of Basal Cell Markers in the Differential Diagnosis of HGPIN

PINATYP

A common scenario where it is difficult to distinguish acinar adenocarcinoma from high grade PIN is when there are a few atypical glands immediately adjacent to high grade PIN. The differential diagnosis is whether these small glands represent tangential sectioning or outpouching off of the high grade PIN glands or a small focus of carcinoma adjacent to the high grade. We refer to these foci as PINATYP. A diagnosis of carcinoma can be rendered only if the small atypical glands are too numerous or too far away from the high grade PIN glands to represent outpouching or tangential sectioning from the PIN glands. In cases of PINATYP, the lack of basal cells in the small atypical glands can be construed as evidence that these glands represent infiltrating cancer only if there are more than a few such glands. As high grade PIN glands can have discontinuous basal cells, one can envision tangential sections off PIN glands in which all cells would appear negative for basal cell markers, such that a few negative small atypical glands adjacent to PIN is not diagnostic of cancer. Some cases may have the appearance of PINATYP yet

will be entirely negative for basal cell markers; these foci may be diagnostic of cancer if there are a sufficiently large number of glands that are not immunoreactive. One may also see classic high grade PIN where some of the glands show the expected patchy basal cell layer and other identical glands are negative for the basal cell markers; these cases we would still diagnose as high grade PIN. Racemase does not differentiate between high grade PIN and cancer, as both typically express this antigen.

Kronz JD, Shaikh AA, Epstein JI. High-grade prostatic intraepithelial neoplasia with adjacent small atypical glands on prostate biopsy. *Hum Pathol* 2001;32:389-95.

Jiang Z, Woda BA, Wu CL, Yang XJ. Discovery and clinical application of a novel prostate cancer marker: alpha-methylacyl CoA racemase (P504S). *Am J Clin Pathol* 2004;122:275-89.

Wu CL, Yang XJ, Tretiakova M, et al. Analysis of alpha-methylacyl-CoA racemase (P504S) expression in high-grade prostatic intraepithelial neoplasia. *Hum Pathol* 2004;35:1008-13.

PIN-like Ductal Adenocarcinoma

A more recently described variant of ductal adenocarcinomas closely resembles high grade prostatic intraepithelial neoplasia (HGPIN) and is composed of simple glands with flat, tufting or micropapillary architecture. PIN-like ductal adenocarcinoma differs from HGPIN by the presence of cystically dilated glands, a greater predominance of flat architecture, and less frequently prominent nucleoli. Verification often requires the immunohistochemical documentation of the absence of basal cells in numerous atypical glands. Although usual ductal adenocarcinoma is considered comparable to Gleason score 8, PIN-like ductal adenocarcinoma was accompanied by Gleason score 6 acinar carcinoma and behaved similar to Gleason score 6 acinar cancer.

Tavora F, Epstein JI. High grade prostatic intraepithelial neoplasia-like ductal adenocarcinoma of the prostate: A clinicopathologic study of 28 cases. *Am J Surg Pathol* (July) 32: 1060,1067, 2008.

Intraductal Carcinoma

Intraductal carcinoma of the prostate (IDC-P) in radical prostatectomy specimens is described as an atypical glandular lesion that spans the entire lumen of prostatic ducts or acini while the normal architecture of ducts or acini is still maintained. Rarely, IDC-P may be identified on biopsy material in the absence of infiltrating carcinoma. Our definition of IDC-P on needle biopsy was derived to identify objective morphological criteria that either architecturally or cytologically clearly exceed those seen in high grade PIN. It is critical to distinguish between high grade PIN and IDC-P, as the former is typically not treated with definitive therapy and recent data has questioned whether high grade PIN on needle biopsy even requires immediate rebiopsy within the first year following its diagnosis. Both entities share cytological features such nuclear enlargement,

hyperchromasia, and enlarged nucleoli. Although dense cribriform (more solid than luminal areas) and solid patterns are not architectural patterns associated with high grade PIN, loose cribriform and micropapillary patterns overlap between the two entities. To establish the diagnosis of IDC-P in the latter two patterns, other cytological features such as markedly enlarged nuclei (6 times larger than those in adjacent non-neoplastic cells) and non-focal comedonecrosis are required. Whereas, it has been accepted that classic high grade PIN can contain a rare gland with focal necrosis, more extensive necrosis is not acceptable. IDC-P also tends to show more prominent nuclear pleomorphism, as opposed to typical high grade PIN with its uniformly enlarged nuclei. Cases which do not satisfy the strict criteria for IDC-P on needle biopsy yet appear more atypical either architecturally or cytologically than usual high grade PIN can be diagnosed as borderline between IDC-P and high grade PIN with a strong recommendation for repeat biopsy.

Infiltrating cribriform acinar adenocarcinoma (Gleason pattern 4 or Gleason pattern 5 with comedonecrosis) closely mimics cribriform IDC-P. Most cases of IDC-P would be diagnosed as cribriform carcinoma if immunohistochemistry demonstrating basal cells had not been performed. In some cases, the contour and branching pattern of normal duct architecture distinguishes IDC-P from infiltrating carcinoma. Ultimately, the presence of a basal cell layer either identified on routine hematoxylin and eosin prepared slides or with immunohistochemistry rules out infiltrating acinar prostate adenocarcinoma. Despite the presence of comedonecrosis, Gleason pattern 5 adenocarcinoma is ruled out also by the identification of a basal cell layer. Although there are extremely rare cases of early small foci of non-cribriform carcinoma of the prostate with focal retention of basal cell layer, this has never been described in cribriform, solid, or micropapillary prostate acinar carcinoma.

McNeal JE, Yemoto CEM. Spread of adenocarcinoma within prostatic ducts and acini. *Am J Surg Pathol* 20:802-814,1996.

Guo CC, Epstein JI. Intraductal carcinoma of the prostate: Histologic features and clinical significance. *Mod Pathol* 19:1528-35, 2006.

Prostate Adenocarcinoma vs. Urothelial Carcinoma

Even in poorly differentiated prostatic carcinomas, there is typically relatively little pleomorphism or mitotic activity compared to poorly differentiated urothelial carcinoma. Poorly differentiated prostate cancers may have enlarged nuclei and prominent nucleoli, yet there is little variability in nuclear shape or size from one nucleus to another. High-grade urothelial carcinomas often reveal marked pleomorphism with tumor giant cells. A subtler finding is that the cytoplasm of prostatic adenocarcinoma is often very foamy and pale imparting a “soft” appearance. In contrast, urothelial carcinomas may demonstrate hard glassy eosinophilic cytoplasm or more prominent squamous differentiation. The findings of infiltrating cords of cells or focal cribriform glandular differentiation are other features more typical of prostatic adenocarcinoma than urothelial carcinoma. Urothelial cancer tends to grow in nests, even when poorly differentiated. Although the above distinction between urothelial carcinoma and prostatic

adenocarcinoma on H&E stained sections is valid for almost all cases, we have seen rare cases where prostate adenocarcinoma has had marked pleomorphism identical to urothelial carcinoma. Consequently, in a poorly differentiated tumor involving the bladder and prostate without any glandular differentiation typical of prostate adenocarcinoma, the case should be worked up immunohistochemically.

Approximately 95% of poorly differentiated prostatic adenocarcinomas show PSA and PSAP staining although it may be focal. While some studies claim superiority of PSA over PSAP in staining prostatic carcinoma, other articles have demonstrated poorly differentiated prostatic carcinomas that lacked PSA staining but still maintained their immunoreactivity with antibodies to PSAP. In our own hands, PSA has in general been more sensitive. Monoclonal antibodies to PSAP have lower sensitivities than their polyclonal counterparts. We have compared PSA staining in a group of poorly differentiated prostatic adenocarcinomas with “poor” PSA staining to newer prostate specific markers including prostate specific membrane antigen (PSMA), p501S (Prostein) and NKX 3.1. Completely negative staining was seen in 15% (PSA), 12% (PSMA), 17% (P501S) and 5% (NKX 3.1) of the cases. Five per cent of the cases were negative for all four markers combined. A similar 5% rate of “false negativity” is found when combining PSA and PSAP stains. Therefore, the lack of immunoreactivity to prostate specific markers in a poorly differentiated tumor within the prostate, especially if present in limited amount, does not exclude the diagnosis of a poorly differentiated prostatic adenocarcinoma.

In a poorly differentiated tumor occurring in the bladder and the prostate where the differential diagnosis is between high-grade prostatic adenocarcinoma and urothelial carcinoma, focal strong staining for either marker can be used reliably to make the diagnosis of prostatic adenocarcinoma, since PSAP and PSA false positivity have not been convincingly described in urothelial carcinomas.

In general, various cytokeratins (CK7, CK20, high molecular weight cytokeratin) show strong positivity in cases of urothelial carcinoma involving the prostate. Although CK7 and CK20 are more frequently seen in urothelial carcinoma as compared to adenocarcinoma of the prostate, they may also be positive in adenocarcinoma of the prostate, such that in our experience they are not that helpful in this differential diagnosis. We and others have found high molecular weight cytokeratin to be positive in more than 90% of urothelial carcinomas. In contrast, high molecular weight cytokeratin is only rarely (8%) expressed, and usually in a very small percentage of cells, in adenocarcinoma of the prostate. P63 is another useful marker in differentiating high grade urothelial from prostatic adenocarcinoma. Using tissue microarrays, we found p63 to have a greater specificity albeit lower sensitivity for urothelial carcinoma compared to high molecular weight cytokeratin (100% specificity and 83% sensitivity). Other markers that also appear highly specific but only of modest sensitivity for urothelial carcinoma include uroplakin and thrombomodulin (49%-69 % sensitivity).

Svanholm H. Evaluation of commercial immunoperoxidase kits for prostatic specific antigen and prostatic specific acid phosphatase. *Acta Pathol Microbiol Immunol Scand [A]* 1986;94:7-12.

Ellis DW, Leffers S, Davies JS, Ng AB. Multiple immunoperoxidase markers in benign

hyperplasia and adenocarcinoma of the prostate. *Am J Clin Pathol* 1984;81:279-84.

Ford TF, Butcher DN, Masters JR, Parkinson MC. Immunocytochemical localisation of prostate-specific antigen: specificity and application to clinical practice. *Br J Urol* 1985;57:50-5.

Feiner HD, Gonzalez R. Carcinoma of the prostate with atypical immunohistological features. Clinical and histologic correlates. *Am J Surg Pathol* 1986;10:765-70.

Keillor JS, Aterman K. The response of poorly differentiated prostatic tumors to staining for prostate specific antigen and prostatic acid phosphatase: a comparative study. *J Urol* 1987;137:894-6.

Epstein JI. PSAP and PSA as immunohistochemical markers. *Urol Clin North Am* 1993;20:757-70.

Chuang A-Y, DeMarzo AM, Veltri RW, Sharma RB, Bieberich CJ and Epstein JI. Immunohistochemical differentiation of high-grade prostate carcinoma from urothelial carcinoma. *Am J Surg Pathol* (in press).

Mhaweck P, Uchida T, Pelte MF. Immunohistochemical profile of high-grade urothelial bladder carcinoma and prostate adenocarcinoma. *Hum Pathol* 2002;33:1136-40.

Nadji M, Tabei SZ, Castro A, et al. Prostatic-specific antigen: an immunohistologic marker for prostatic neoplasms. *Cancer* 1981;48:1229-32.

Genega EM, Hutchinson B, Reuter VE, Gaudin PB. Immunophenotype of high-grade prostatic adenocarcinoma and urothelial carcinoma. *Mod Pathol* 2000;13:1186-91.

Varma M, Morgan M, Amin MB, Wozniak S, Jasani B. High molecular weight cytokeratin antibody (clone 34betaE12): a sensitive marker for differentiation of high-grade invasive urothelial carcinoma from prostate cancer. *Histopathology* 2003;42:167-72.